GUIDELINES FOR EXTRACTION CONSIDERING BLOOD PRESSURE IN ADULTS

OUTLINE

Introduction
Guidelines by ASA
Conclusion

Introduction:

Management of Hypertensive Patients in a Dental Clinic: Proposed Guidelines by the American Society of Anesthesiologists (ASA)

Guidelines by ASA:

BLOOD PRESSURE	ASA CLASSIFICATION	DENTAL THERAPY CONSIDERATIONS
<140 and<90	Ι	 Pay attention to routine dental care Conduct another examination after six months
140 -159 and/ or 90-94	Π	 Before receiving dental treatment, recheck blood pressure three times in a row. If all readings are higher than prescribed levels, a trip to the doctor is advised. Pay attention to standard dental care. Follow the recommended stress-reduction strategy.
160-199 and/or 95-144	III	• Check blood pressure again in five minutes.

DEPARTMENT OF ORAL AND MAXILLOFACIAL SURGERY

BLOOD PRESSURE	ASA CLASSIFICATION	DENTAL THERAPY CONSIDERATIONS
		 If the condition persists, get medical advice before starting dental treatment. Pay attention to routine dental care Put a stress-reduction plan into action.
>200 and /or >115	IV	 Check blood pressure again in five minutes. If blood pressure is increased, avoid doing dental therapy, either on a regular basis or in an emergency situation. Avoid dental therapy if blood pressure is elevated, whether it be routinely or in an emergency.

CONCLUSION

Therefore, if a hypertensive patient enters a dental clinic, the instructions should be followed.

Reference:

* Malamed, Stanley. Medical Emergencies in the Dental Office - E-Book. 7th ed. Elsevier Health Sciences, 2014. Web.15 OCT.2022.

Author: Dr .SaneemAhamed.A.